



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MISCELLANEOUS.

130. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, Ohio.

From a cloud of angular elevation $\phi=45^\circ$, a streak of lightning darted to the earth. The temperature of the atmosphere was $t=80^\circ$, and the percentage of humidity $p=90$. After $m=3$ seconds, the report of the stroke at the earth was heard. How far away from the observer did the streak of lightning (1) start, and (2) strike the earth?

Solution by G. B. M. ZERR, A. M., Ph.D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Let H =tension of aqueous vapor at $t^\circ=80^\circ$.

Let H' =tension of aqueous vapor at dew point.

Then $(H'/H)100=p=90$, $80^\circ F=26^\circ.7 C$.

Now $H=26.045\text{mm}$. $\therefore H'=\frac{9}{10}$ of $26.045\text{mm}=23.4405\text{mm}$.

Let us take the barometric pressure at $740\text{mm}=P$.

Then $740\text{mm}-23.4405\text{mm}=716.5595\text{mm}$. will represent the pressure of dry air.

Let W =weight of litre of dry air at $80^\circ F$.

W' =weight of vapor in one litre.

$$W=1.293187 \times \frac{716.5595}{760} \times \frac{1}{1+.00367 \times 26.7}=1.110667 \text{ grams.}$$

$$W'=1.293187 \times .6235 \times \frac{23.4405}{760} \times \frac{1}{1+.00367 \times 26.7}=.022653 \text{ grams.}$$

$W+W'=1.13332$ grams=total weight of litre.

$V=\sqrt{(Pk/\rho)}$, where $k=1.41$ =ratio of specific heat of air at constant pressure to the specific heat at constant volume.

$P=740\text{mm}=986634$ dynes per square cm., ρ =density=.00113332.

$\therefore V=\sqrt{[(986634 \times 1.41)/.00113332]}=350.36$ meters per second.

$3 \times 60 \times 350.36=63.0648$ kilometers=distance of observer from point on earth where lightning strikes. If we regard the distance as a straight line, then distance to cloud= $63.0648/\sqrt{2}=89.18624$ kilometers.

If we regard the earth as a sphere, diameter 6370.946 kilometers, then $1^\circ=111\frac{1}{3}$ kilometers. Therefore $34^\circ 3' 3''=63.0648$ kilometers.

$$\text{Distance to cloud}=\frac{6370.946 \sin(34^\circ 3' 3'')}{\sin(44^\circ 25' 56.7'')}=91.4594 \text{ kilometers.}$$

131. Proposed by SAUL EPSTEIN, Ph. D., The University of Chicago.

Find a power series for π^{n^x} (n =any integer).

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

$$\pi^{n^x}=1+n x \log_e \pi + \frac{n^2 x^2}{2!} (\log_e \pi)^2 + \frac{n^3 x^3}{3!} (\log_e \pi)^3 + \dots$$